

TITLE OF THE INVENTION

BUCCAL, POLAR AND NON-POLAR SPRAY OR CAPSULE

RELATED APPLICATIONS

- 5 This application is a continuation in part of applicant PCT application PCT/US97/17899 filed October 1st 1997.

BACKGROUND OF THE INVENTION

- 10 It is known that certain biologically active compounds are better absorbed through the oral mucosa than through other routes of administration, such as through the stomach or intestine. However, formulations suitable for such administration by these latter routes present their own problems. For example, the biologically active compound must be compatible with the other components of the composition such as
- 15 propellants, solvents, etc. Many such formulations have been proposed. For example, U.S.P. 4,689,233, Dvorsky et al., describes a soft gelatin capsule for the administration of the anti-coronary drug nifedipine dissolved in a mixture of polyether alcohols. U.S.P. 4,755,389, Jones et al., describes a hard gelatin chewable capsule containing nifedipine. A
- 20 chewable gelatin capsule containing a solution or dispersion of a drug is described in U.S.P. 4,935,243, Borkan et al. U.S.P. 4,919,919, Aouda et al. and U.S.P. 5,370,862, Klokke-Bethke, describe a nitroglycerin spray for administration to the oral mucosa comprising nitroglycerin, ethanol, and other components. An orally administered pump spray is described by
- 25 Cholcha in U.S.P. 5,186,925. Aerosol compositions containing a hydrocarbon propellant and a drug for administration to a mucosal surface are described in U.K. 2,082,457, Su, U.S.P. 3,155,574, Silson et al., U.S.P. 5,011,678, Wang et al., and by Parnell in U.S.P. 5,128,132. It should be noted that these references discuss bioavailability of solutions by
- 30 inhalation rather than through the membranes to which they are administered.

SUMMARY OF THE INVENTION

A buccal aerosol spray or soft bite gelatin capsule using a polar or non-polar solvent has now been developed which provides biologically active compounds for rapid absorption through the oral mucosa, resulting in fast onset of effect.

The buccal aerosol spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable non-polar solvent comprise in weight % of total composition: pharmaceutically acceptable propellant 5-80%, non-polar solvent 20-85%, active compound 0.05-50%, suitably additionally comprising, by weight of total composition a flavoring agent 0.01-10%. Preferably the composition comprises: propellant 10-85%, non-polar solvent 25-89.9%, active compound 0.01-40%, flavoring agent 1-8%; most suitably propellant 20-70%, non-polar solvent 30-74.75%, active compound 0.25-35%, flavoring agent 2-7.5%.

The buccal polar aerosol spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable polar solvent are also administrable in aerosol form driven by a propellant. In this case the composition comprise in weight% of total composition: aqueous polar solvent 10-99%, active compound 0.1-25%, suitably additionally comprising, by weight of total composition a flavoring agent 0.05-10% and propellant: 2 - 10%. Preferably the composition comprises: polar solvent 20 - 97%, active compound 0.1-15%, flavoring agent 0.1-5% and propellant: 3 - 5%; most suitably polar solvent 25 - 97%, active compound 0.2-25%, flavoring agent 0.1-2.5% and propellant: 3 - 4%.

The buccal pump spray composition of the present invention for transmucosal administration of a pharmacologically active compound where

said active compound is soluble in a pharmacologically acceptable non-polar solvent said composition comprise in weight % of total composition: non-polar solvent 30-99.69%, active compound 0.005-55%, and suitably additionally, flavoring agent 0.1-10%.

The buccal polar pump spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable polar solvent comprising in weight% of total composition: aqueous polar solvent 30-99.69%, active compound 0.001-60%, suitably additionally comprising, by weight of total composition a flavoring agent 0.1-10%. Preferably the composition comprises: polar solvent 37-98.58%, active compound 0.005-55%, flavoring agent 0.5-8%; most suitably polar solvent 60.9-97.06%, active compound 0.01-40%, flavoring agent 0.75-7.5%.

The soft bite gelatin capsules of the present invention for transmucosal administration of a pharmacologically active compound, at least partially soluble in a pharmacologically acceptable non-polar solvent, having charged thereto a fill composition comprise in weight % of total composition: non-polar solvent 4-99.99%, emulsifier 0-20%, active compound 0.01-80%, provided that said fill composition contains less than 10% of water, suitably additionally comprising, by weight of the composition: flavoring agent 0.01-10%. Preferably, the soft bite gelatin capsule comprises: non-polar solvent 21.5-99.975%, emulsifier 0-15%, active compound 0.025-70%, flavoring agent 1-8%; most suitably: non-polar solvent 28.5-97.9%, emulsifier 0-10%, active compound 0.1-65.0%, flavoring agent 2-6%.

The soft bite polar gelatin capsules of the present invention for transmucosal administration of a pharmacologically active compound, at least partially soluble in a pharmacologically acceptable polar solvent, having charged thereto a composition comprising in weight % of total composition:

006220-8112500

polar solvent 25-99.89%, emulsifier 0-20%, active compound 0.01-65%, provided that said composition contains less than 10% of water, suitably additionally comprising, by weight of the composition: flavoring agent 01-10%. Preferably, the soft bite gelatin capsule comprises: polar solvent 5 37-99.95%, emulsifier 0-15%, active compound 0.025-55%, flavoring agent 1-8%; most suitably: polar solvent 44-96.925%, emulsifier 0-10%, active compound 0.075-50%, flavoring agent 2-6%.

10 It is an object of the invention to coat the mucosal membranes either with extremely fine droplets of spray containing the active compounds or a solution or paste thereof from bite capsules.

15 It is also an object of the invention to administer to the oral mucosa of a mammalian in need of same, preferably man, by spray or bite capsule. a predetermined amount of a biologically active compound by this method or from a soft gelatin bite capsule.

20 A further object is a sealed aerosol spray container containing a composition of the non polar or polar aerosol spray formulation, and a metered valve suitable for releasing from said container a predetermined amount of said composition.

25 As the propellant evaporates after activation of the aerosol valve, a mist of fine droplets is formed which contains solvent and active compound.

30 The propellant is a non-Freon material, preferably a C_{3-8} hydrocarbon of a linear or branched configuration. The propellant should be substantially non-aqueous. The propellant produces a pressure in the aerosol container such that under expected normal usage it will produce sufficient pressure to expel the solvent from the container when the valve is activated but not excessive pressure such as to damage the container or valve seals.

09537118-032900

The non-polar solvent is a non-polar hydrocarbon, preferably a C₇₋₁₈ hydrocarbon of a linear or branched configuration, fatty acid esters, and triglycerides, such as miglyol. The solvent must dissolve the active compound and be miscible with the propellant, i.e., solvent and propellant must form a single phase at 0-40°C at a pressure range of 1-3 atm.

The polar and non-polar aerosol spray compositions of the invention are intended to be administered from a sealed, pressurized container. Unlike a pump spray, which allows the entry of air into the container after every activation, the aerosol container of the invention is sealed at the time of manufacture. The contents of the container are released by activation of a metered valve, will does not allow entry of atmospheric gasses with each activation. Such containers are commercially available.

A further object is a pump spray container containing a composition of the pump spray formulation, and a metered valve suitable for releasing from said container a predetermined amount of said composition.

A further object is a soft gelatin bite capsule containing a composition of as set forth above. The formulation may be in the form of a viscous solution or paste containing the active compounds. Although solutions are preferred, paste fills may also be used where the active compound is not soluble or only partially soluble in the solvent of choice. Where water is used to form part of the paste composition, it should not exceed 10% thereof. (All percentages herein are by weight unless otherwise indicated.)

The polar or non-polar solvent is chosen such that it is compatible with the gelatin shell and the active compound. The solvent preferably dissolves the active compound. However, other components wherein the active compound is not soluble or only slightly soluble may be used and will form a paste fill.

Soft gelatin capsules are well known in the art. See, for example, U.S.P. 4,935,243, Borkan et al., for its teaching of such capsules. The capsules of the present invention are intended to be bitten into to release the low viscosity solution or paste therein, which will then coat the buccal mucosa with the active compounds. Typical capsules, which are swallowed whole or bitten and then swallowed, deliver the active compounds the stomach, which results in significant lag time before maximum blood levels can be achieved or subject the compound to a large first pass effect. Because of the enhanced absorption of the compounds through the oral mucosa and no chance of a first pass effect, use of the bite capsules of the invention will eliminate much of the lag time, resulting in hastened onset of biological effect. The shell of a soft gelatin capsule of the invention may comprise, for example:

- 15 Gelatin: 50-75%, glycerin 20-30%, colorants 0.5-1.5%, water 5-10%, and sorbitol 2-10%.

The active compound may include biologically active peptides, central nervous system active amines, sulfonyl ureas, antibiotics, antifungals, anti-virals, sleep inducers, antiasthmatics, bronchial dilators, antiemetics, histamine H-2 receptor antagonists, barbiturates, prostaglandins and neutraceuticals.

The active compounds may also include antihistamines, alkaloids, hormones, benzodiazepines and narcotic analgesics. While not limited thereto, these active compounds are particularly suitable for non-polar pump spray formulation and application.

BRIEF DESCRIPTION OF THE DRAWING

30 The figure is a schematic diagram showing routes of absorption and processing of pharmacologically active substances in a mammalian system.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

0053718-032000

5 The preferred active compounds of the present invention are in an ionized, salt form or as the free base of the pharmaceutically acceptable salts thereof (provided, for the aerosol or spray compositions, they are soluble in the spray solvent). These compounds are soluble in the non-polar solvents of the invention at useful concentrations or can be prepared as pastes at useful concentrations. These concentrations may be less than the standard accepted dose for these compounds since there is enhanced absorption of the compounds through the oral mucosa. This aspect of the
10 invention is especially important when there is a large (40-99.99%) First pass effect.

As propellants for the non polar sprays, propane, N-butane, iso-butane, N-pentane, iso-pentane, and neo-pentane, and mixtures thereof may be
15 used. N-butane and iso-butane, as single gases, are the preferred propellants. It is permissible for the propellant to have a water content of no more than 0.2%, typically 0.1-0.2%. (All percentages herein are by weight unless otherwise indicated.) It is also preferable that the propellant be synthetically produced to minimize the presence of contaminants which
20 are harmful to the active compounds. These contaminants include oxidizing agents, reducing agents, Lewis acids or bases, and water. The concentration of each of these should be less than 0.1%, except that water may be as high as 0.2%.

25 Suitable non-polar solvents for the capsules and the non-polar sprays include (C₂-C₂₄) fatty acid C₂-C₆ esters, C₇-C₁₈ hydrocarbon, C₂-C₆ alkanoyl esters, and the triglycerides of the corresponding acids. When the capsule fill is a paste, other liquid components may be used instead of the above low molecular weight solvents. These include soya oil, corn oil, other
30 vegetable oils.

As solvents for the polar capsules or sprays there may be used low molecular weight polyethyleneglycols (PEG) of 400-1000 Mw (preferably 400-600), low molecular weight (C_2-C_8) mono and polyols and alcohols of C_7-C_{18} linear or branch chain hydrocarbons, glycerin may also be present and water may also be used in the sprays, but only in limited amount in the capsules.

It is expected that some glycerin and water used to make the gelatin shell will migrate from the shell to the fill during the curing of the shell.

Likewise, there may be some migration of components from the fill to the shell during curing and even throughout the shelf-life of the capsule. Therefore, the values given herein are for the compositions as prepared, it being within the scope of the invention that minor variations will occur.

The preferred flavoring agents are synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, sweeteners (sugars, aspartame, saccharin, etc.), and combinations thereof.

The active substances include the active compounds selected from the group consisting of cyclosporine, sermorelin, Octreotide acetate, calcitonin-salmon, insulin lispro, sumatriptan succinate, clozapine, cyclobenzaprine, dexfenfluramine hydrochloride, glyburide, zidovudine, erythromycin, ciprofloxacin, ondansetron hydrochloride, dimenhydrinate, cimetidine hydrochloride, famotidine, phenytoin sodium, phenytoin, carboprost tromethamine, carboprost, diphenhydramine hydrochloride, isoproterenol hydrochloride, terbutaline sulfate, terbutaline, theophylline, albuterol sulfate and neutraceuticals, that is to say nutrients with pharmacological action such as but not limited to carnitine, valerian, echinacea, and the like.

The formulations of the present invention comprise an active compound or a pharmaceutically acceptable salt thereof. The term "pharmaceutically acceptable salts" refers to salts prepared from

pharmaceutically acceptable non-toxic acids or bases including organic and inorganic acids or bases.

006220 BT 2560

When an active compound of the present invention is acidic, salts
5 may be prepared from pharmaceutically acceptable non-toxic bases. Salts
derived from all stable forms of inorganic bases include aluminum,
ammonium, calcium, copper, iron, lithium, magnesium, manganese,
potassium, sodium, zinc, etc. Particularly preferred are the ammonium,
calcium, magnesium, potassium, and sodium salts. Salts derived from
10 pharmaceutically acceptable organic non-toxic bases include salts of
primary, secondary, and tertiary amines, substituted amines including
naturally occurring substituted amines, cyclic amines and basic ion-
exchange resins such as arginine, betaine, caffeine, choline, N,N'-
dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethyl-
15 aminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethyl-
piperidine, glucamine, glucosamine, histidine, isopropylamine, lysine,
methyl-glucosamine, morpholine, piperazine, piperidine, polyamine resins,
procaine, purine, theobromine, triethylamine, trimethylamine, tripropylamine,
etc.

20

When an active compound of the present invention is basic, salts
may be prepared from pharmaceutically acceptable non-toxic acids. Such
acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric,
ethane-sulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric,
25 isethionic, lactic, maleic, mandelic, methanesulfonic, mucic, nitric, pamoic,
pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic, etc.
Particularly preferred are citric, hydrobromic, maleic, phosphoric, sulfuric,
and tartaric acids.

30

In the discussion of methods of treatment herein, reference to the
active compounds is meant to also include the pharmaceutically acceptable
salts thereof. While certain formulations are set forth herein, the actual

000322 RV1

10

PHCO3.0-008CIP

amounts to be administered to the mammal or man in need of same are to be determined by the treating physician.

The invention is further defined by reference to the following
5 examples, which are intended to be illustrative and not limiting.

000000 000000 000000

The following are examples of each class (all values unless otherwise specified are in weight percent):

EXAMPLE 1

5

Biologically active peptides including peptide hormones

A. Cyclosporine lingual spray

	Amounts	preferred amount	most preferred amount
Cyclosporine	5-50	10-35	15-25
water	5-20	7.5-50	9.5-12
ethanol	5-60	7.5-50	10-20
polyethylene glycol	20-60	30-45	35-40
flavors	0.1-5	1-4	2-3

Sub
B. 7B. Cyclosporine Non-Polar lingual spray

	Amounts	preferred amount	most preferred amount
Cyclosporine	1-50	3-40	5-30
Migylol	20-70	25-1	30-40
Polyoxyethyl- ated castor oil	20 -	25 -1	30-40
Butane	25-80	30-70	33-50
flavors	0.1-5	1-4	2-3

09537118-032800

C. Cyclosporine non-polar bite capsule

	Amounts	preferred amount	most preferred amount
Cyclosporine	1-35	5-25	10-20
olive oil	25-60	35-55	30-45
polyoxyethyl- ated oleic glycerides	25-60	35-55	30-45
flavors	0.1-5	1-4	2-3

D. Cyclosporine bite capsule

	Amounts	preferred amount	most preferred amount
5 Cyclosporine	5-50	10-35	15-25
polyethylene glycol	20-60	30-45	35-40
glycerin	5-30	7.5-25	10-20
propylene glycol	5-30	7.5-25	10-20
flavors	0.1-10	1-8	3-6

10

E. Sermorelin (as the acetate) lingual spray

	Amounts	preferred amount	most preferred
sermorelin (as the acetate)	.01-5	.1-3	.2-1.0
mannitol,	1-25	5-20	10-15
15 monobasic sodium phosphate,	0.1-5	1-3	1.5-2.5
dibasic sodium phosphate water	0.01-5	.05-3	0.1-0.5
ethanol	5-30	7.5-25	9.5-15
polyethylene glycol	20-60	30-45	35-40
propylene glycol	5-25	10-20	12-17
20 flavors	0.1-5	1-4	2-3

000322 RV1 000322 RV1

F. Octreotide acetate (Sandostatin®) lingual spray

	Amounts	preferred amount	most preferred amount
octreotide acetate	0.001-0.5	0.005-0.250	0.01-0.10
acetic acid	1-10	2-8	4-6
5 sodium acetate	1-10	2-8	4-6
sodium chloride	3-30	5-25	15-20
flavors	0.1-5	0.5-.4	2-3
ethanol	5-30	7.5-20	9.5-15
water	15-95	35-90	65-85
10 flavors	0.1-5	1-4	2-3

G. Calcitonin-salmon lingual spray

	Amounts	preferred amount	most preferred amount
Calcitonin-salmon	0.001-5	0.005-2	01-1.5
15 ethanol	2-15	3-10	7-9.5
water	30-95	50-90	60-80
polyethylene glycol	2-15	3-10	7-9.5
sodium chloride	2.5-20	5-15	10-12.5
flavors	0.1-5	1-4	2-3

H. insulin lispro, lingual spray

	Amounts	preferred amount	most preferred amount
insulin,	20-60	4-55	5-50
glycerin,	0.1-10	0.25-5	0.1-1.5
25 dibasic sodium phosphate,	1-15	2.5-10	4-8
m-cresol,	1-25	5-25	7.5-12.5
zinc oxide	0.01-0.25	.05-0.15	0.075-0.10
m-cresol,	0.1-1	0.2-0.8	0.4-0.6
phenol	trace amounts	trace amounts	trace amounts
30 ethanol	5-20	7.5-15	9-12
water	30-90	40-80	50-75
propylene glycol	5-20	7.5-15	9-12
flavors	0.1-5	0.5-3	0.75-2
adjust pH to 7.0-7.8 with HCl or NaOH			

000322 RV1 000322 RV1

EXAMPLE 2

CNS active amines and their salts: including but not limited to tricyclic amines, GABA analogues, thiazides, phenothiazine derivatives, Serotonin antagonists and serotonin reuptake inhibitors

00537119-032000

5

A. Sumatriptan succinate lingual spray

	Amounts	preferred amount	most preferred amount
sumatriptan succinate	0.5-30	1-20	10-15
ethanol	5-60	7.5-50	10-20
propylene glycol	5-30	7.5-20	10-15
polyethylene glycol	0-60	30-45	35-40
water	5-30	7.5-20	10-15
flavors	0.1-5	1-4	2-3

10

B. Sumatriptan succinate bite capsule

	Amounts	preferred amount	most preferred amount
sumatriptan succinate	0.01-5	0.05-3.5	0.075-1.75
polyethylene glycol	25-70	30-60	35-50
glycerin	25-70	30-60	35-50
flavors	0.1-10	1-8	3-6

15

20

C. Clozepine lingual spray

	Amounts	preferred amount	most preferred amount
Clozepine	0.5-30	1-20	10-15
ethanol	5-60	7.5-50	10-20
propylene glycol	5-30	7.5-20	10-15
polyethylene glycol	0-60	30-45	35-40
water	5-30	7.5-20	10-15
flavors	0.1-5	1-4	2-3

25

D. Clozepine Non-Polar lingual spray with propellant

	Amounts	preferred amount	most preferred amount
Clozepine	0.5-30	1-20	10-15
Migylol	20-85	25-70	30-40
Butane	15-80	30-75	60-70
flavors	0.1-5	1-4	2-3

E. Clozepine Non-Polar lingual spray without propellant

	Amounts	preferred amount	most preferred amount
Clozepine	0.5-30	1-20	10-15
Migylol	70-99.5	80-99	85-90
flavors	0.1-5	1-4	2-3

5

F. Cyclobenzaprine Non polar lingual spray

	Amounts	preferred amount	most preferred amount
Cyclobenzaprine (base)	0.5-30	1-20	10-15
Migylol	20-85	25-70	30-40
Iso-butane	15-80	30-75	60-70
flavors	0.1-5	1-4	2-3

G. dexfenfluramine hydrochloride lingual spray

	Amounts	preferred amount	most preferred amount
10 dexfenfluramine Hcl	5-30	7.5-20	10-15
ethanol	5-60	7.5-50	10-20
propylene glycol	5-30	7.5-20	10-15
polyethylene glycol	0-60	30-45	35-40
water	5-30	7.5-20	10-15
15 flavors	0.1-5	1-4	2-3

000322 RV1 000322 RV1

EXAMPLE 3

Sulfonylureas

A. Glyburide lingual spray

		Amounts	preferred amount	most preferred amount
5	Glyburide	0.25-25	0.5-20	0.75-15
	ethanol	5-60	7.5-50	10-20
	propylene glycol	5-30	7.5-20	10-15
	polyethylene glycol	0-60	30-45	35-40
	water	2.5-30	5-20	6-15
10	flavors	0.1-5	1-4	2-3

B. Glyburide non-polar bite capsule

		Amounts	preferred amount	most preferred amount
	Glyburide	0.01-10	0.025-7.5	0.1-4
	olive oil	30-60	35-55	30-50
	polyoxyethyl- ated oleic glycerides	30-60	35-55	30-50
	flavors	0.1-5	1-4	2-3

EXAMPLE 4

Antibiotics anti-fungals and anti-virals

15 A. zidovudine [formerly called azidothymidine (AZT) (Retrovir) non-polar lingual

		Amounts	spray preferred amount	most preferred amount
	zidovudine	10-50	15-40	25-35
	Soya oil	20-85	25-70	30-40
	Butane	15-80	30-75	60-70
	flavors	0.1-5	1-4	2-3

000322 RV1-032200

B. Erythromycin bite capsule

	Amounts	preferred amount	most preferred amount
Erythromycin	25-65	30-50	35-45
polyoxyethylene glycol	5-70	30-60	45-55
5 glycerin	5-20	7.5-15	10-12.5
flavors	1-10	2-8	3-6

C. Ciprofloxacin hydrochloride bite capsule

	Amounts	preferred amount	most preferred amount
10 Ciprofloxacin hydrochloride	25-65	35-55	40-50
glycerin	5-20	7.5-15	10-12.5
polyethylene glycol	20-75	30-65	40-60
flavors	1-10	2-8	3-6

D. zidovudine [formerly called azidothymidine (AZT) (Retrovir) lingual spray

	Amounts	preferred amount	most preferred amount
zidovudine	10-50	15-40	25-35
water	30-80	40-75	45-70
20 ethanol	5-20	7.5-15	9.5-12.5
polyethylene glycol	5-20	7.5-15	9.5-12.5
flavors	0.1-5	1-4	2-3

EXAMPLE 5

Anti-emetics

A. Ondansetron hydrochloride lingual spray

	Amounts	preferred amount	most preferred amount
ondansetron hydrochloride	1-25	2-20	2.5-15
citric acid monohydrate,	1-10	2-8	2.5-5
30 sodium citrate dihydrate	0.5-5	1-4	1.25-2.5
water	1-90	5-85	10-75
ethanol	5-30	7.5-20	9.5-15
propylene glycol	5-30	7.5-20	9.5-15
polyethylene glycol	5-30	7.5-20	9.5-15
35 flavors	1-10	3-8	5-7.5

000322 RV1

18

PHCO3.0-008CIP

B. Dimenhydrinate bite capsule

	Amounts	preferred amount	most preferred amount
Dimenhydrinate	0.5-30	2-25	3-15
glycerin	5-20	7.5-15	10-12.5
polyethylene glycol	45-95	50-90	55-85
flavors	1-10	2-8	3-6

C. Dimenhydrinate polar lingual spray

	Amounts	preferred amount	most preferred amount
Dimenhydrinate	3-50	4-40	5-35
water	5-90	10-80	15-75
ethanol	1-80	3-50	5-10
polyethylene glycol	1-80	3-50	5-15
Sorbitol	0.1-5	0.2-4	0.4-1.0
aspartame	0.01-0.5	0.02-0.4	0.04-0.1
flavors	0.1-5	1-4	2-3

10

EXAMPLE 6**Histamine H-2 receptor antagonists****A. Cimetidine hydrochloride bite capsule**

	Amounts	preferred amount	most preferred amount
Cimetidine Hcl	10-60	15-55	25-50
glycerin	5-20	7.5-15	10-12.5
polyethylene glycol	20-90	25-85	30-75
flavors	1-10	2-8	3-6

B. Famotidine lingual spray

	Amounts	preferred amount	most preferred amount
Famotidine	1-35	5-30	7-20
water	2.5-25	3-20	5-10
L-aspartic acid	0.1-20	1-15	5-10
polyethylene glycol	20-97	30-95	50-85
flavors	0.1-10	1-7.5	2-5

C. Famotidine non-polar lingual spray

	Amounts	preferred amount	most preferred amount
Famotidine	1-35	5-30	7-20
Soya oil	10-50	15-40	15-20
Butane	15-80	30-75	45-70
polyoxyethyl- ated oleic glycerides	10-50	15-40	15-20
flavors	0.1-5	1-4	2-3

EXAMPLE 7

Barbiturates

5 A. Phenytoin sodium lingual spray

	Amounts	preferred amount	most preferred amount
Phenytoin sodium	10-60	15-55	20-40
water	2.5-25	3-20	5-10
ethanol	5-30	7.5-20	9.5-15
10 propylene glycol	5-30	7.5-20	9.5-15
polyethylene glycol	5-30	7.5-20	9.5-15
flavors	1-10	3-8	5-7.5

B. Phenytoin non-polar lingual spray

	Amounts	preferred amount	most preferred amount
Phenytoin	5-45	10-40	15-35
migylol	10-50	15-40	15-20
Butane	15-80	30-75	60-70
polyoxyethyl- ated oleic glycerides	10-50	15-40	15-20
flavors	0.1-10	1-8	5-7.5

EXAMPLE 8

Prostaglandins

A. Carboprost thromethamine lingual spray

	Amounts	preferred amount	most preferred amount
5 Carboprost thromethamine	0.05-5	0.1-3	0.25-2.5
water	50-95	60-80	65-75
ethanol	5-20	7.5-15	9.5-12.5
polyethylene glycol	5-20	7.5-15	9.5-12.5
sodium chloride	1-20	3-15	4-8
10 flavors	0.1-5	1-4	2-3

pH is adjusted with sodium hydroxide and/or hydrochloric acid

B. Carboprost non-polar lingual spray

	Amounts	preferred amount	most preferred amount
Carboprost	0.05-5	0.1-3	0.25-2.5
miglyol	25-50	30-45	35-40
Butane	5-60	10-50	20-35
polyoxyethylated oleic glycerides	25-50	30-45	35-40
flavors	0.1-10	1-8	5-7.5

15

EXAMPLE 9

Neutraceuticals

A. Carnitine as bite capsule (contents are a paste)

	Amounts	preferred amount	most preferred amount
20 Carnitine fumarate	6-80	30-70	45-65
soya oil	7.5-50	10-40	12.5-35
soya lecithin	0.001-1.0	0.005-0.5	.01-0.1
Soya fats	7.5-50	10-40	12.5-35
flavors	1-10	2-8	3-6

000322 RV1

EXAMPLE 10

Sleep Inducers (also CNS active amine)

A. Diphenhydramine hydrochloride lingual spray

	Amounts	preferred amount	most preferred amount
Diphenhydramine Hcl	3-50	4-40	5-35
water	5-90	10-80	50-75
ethanol	1-80	3-50	5-10
polyethylene glycol	1-80	3-50	5-15
Sorbitol	0.1-5	0.2-4	0.4-1.0
aspartame	0.01-0.5	0.02-0.4	0.04-0.1
flavors	0.1-5	1-4	2-3

5

EXAMPLE 11

Anti-Asthmatics-Bronchodilators

A. Isoproterenol Hydrochloride as polar lingual spray

	Amounts	preferred amount	most preferred amount
Isoproterenol Hydrochloride	0.1-10	0.2-7.5	0.5-6
water	5-90	10-80	50-75
ethanol	1-80	3-50	5-10
polyethylene glycol	1-80	3-50	5-15
Sorbitol	0.1-5	0.2-4	0.4-1.0
aspartame	0.01-0.5	0.02-0.4	0.04-0.1
flavors	0.1-5	1-4	2-3

00537110:0322000

B. Terbutaline sulfate as polar lingual spray

	Amounts	preferred amount	most preferred amount
Terbutaline sulfate	0.1-10	0.2-7.5	0.5-6
water	5-90	10-80	50-75
ethanol	1-10	2-8	2.5-5
Sorbitol	0.1-5	0.2-4	0.4-1.0
aspartame	0.01-0.5	0.02-0.4	0.04-0.1
flavors	0.1-5	1-4	2-3

C. Terbutaline as non-polar lingual spray

	Amounts	preferred amount	most preferred amount
Terbutaline	0.1-10	0.2-7.5	0.5-6
miglyol	25-50	30-45	35-40
isobutane	5-60	10-50	20-35
polyoxyethylated oleic glycerides	25-50	30-45	35-40
flavors	0.1-10	1-8	5-7.5

5

D. Theophylline polar bite capsule

	Amounts	preferred amount	most preferred amount
Theophylline	5-50	10-40	15-30
polyethylene glycol	20-60	25-50	30-40
glycerin	25-50	35-45	30-40
propylene glycol	25-50	35-45	30-40
flavors	0.1-5	1-4	2-3

000322 RV1-0322000

E. Albuterol sulfate as polar lingual spray

	Amounts	preferred amount	most preferred amount
Albuterol sulfate	0.1-10	0.2-7.5	0.5-6
water	5-90	10-80	50-75
ethanol	1-10	2-8	2.5-5
Sorbitol	0.1-5	0.2-4	0.4-1.0
aspartame	0.01-0.5	0.02-0.4	0.04-0.1
flavors	0.1-5	1-4	2-3

Example 12**Polar solvent formulations using a propellant:**

5

A. Sulfonylurea

	Amount	Preferred Amount	Most-Preferred Amount
Glyburide	0.1-25%	0.5-15%	0.6-10%
Ethanol	40-99%	60-97%	70-97%
Water	0.01-5%	0.1-4%	0.2-2%
Flavors	0.05-10%	0.1-5%	0.1-2.5%
Propellant	2-10%	3-5%	3-4%

10 **B. Prostaglandin E₁ (vasodilator)**

	Amount	Preferred Amount	Most-Preferred Amount
Prostaglandin E ₁	0.01-10%	0.1-5%	0.2-3%
Ethanol	10-90%	20-75%	25-50%
Propylene glycol	1-90%	5-80%	10-75%
Water	0.01-5%	0.1-4%	0.2-2%
Flavors	0.05-10%	0.1-5%	0.1-2.5%
Propellant	2-10%	3-5%	3-4%

00537 118 032000

C. Promethazine (antiemetic, sleep inducer, and CNS active amine)

	Amount	Preferred Amount	Most-Preferred Amount
Promethazine	1-25%	3-15%	5-12%
Ethanol	10-90%	20-75%	25-50%
Propylene glycol	1-90%	5-80%	10-75%
Water	0.01-5%	0.1-4%	0.2-2%
Flavors	0.05-10%	0.1-5%	0.1-2.5%
Propellant	2-10%	3-5%	3-4%

5

D. Meclizine

	Amount	Preferred Amount	Most-Preferred Amount
Meclizine	1-25%	3-15%	5-12%
Ethanol	1-15%	2-10%	3-6
Propylene glycol	20-98%	5-90%	10-85%
Water	0.01-5%	0.1-4%	0.2-2%
Flavors	0.05-10%	0.1-5%	0.1-2.5%
Propellant	2-10%	3-5%	3-4%

00537118-032000